The Plant Journal (2000) **21**(6), 589–592

Author index

Bacanamwo, M. *see* Freyermuth, S.K.
Barbier-Brygoo, H. *see* Geelen, D.
Baulcombe, D.C. *see* Bendahmane, A.
Bäumlein, H. *see* Reidt, W.
Beck, C.F. *see* Schroda, M.
Becker, J., Kempf, R., Jeblick, W. and Kauss, H. Induction of competence for elicitation of defense responses in cucumber hypocotyls requires proteasome activity 311
Bendahmane, A., Querci, M., Kanyuka, K. and Baulcombe, D.C. *Agrobacterium* transient expression system as a tool for the isolation of disease resistance genes: application to the *Rx2* locus in potato 73
Berbier-Brygoo, H. *see* Frachisse, J.-M.
Beynon, J.L. *see* Bittner-Eddy, P.D.
Bisseling, T. *see* Goedhart, J.
Bittner-Eddy, P.D., Crute, I.R., Holub, E.B. and Beynon, J.L. *RPP13* is a simple locus in *Arabidopsis thaliana* for alleles that specifiy downy mildew resistance to different avirulence determinants in *Peronospora parasitica* 177
Bleecker, A.B. *see* Grbić, V.
Blöcker, D. *see* Schroda, M.
Bonin, C.P. and Reiter, W.-D. A bifunctional epimerase-reductase acts downstream of the *MUR1* gene product and completes the *de novo* synthesis of GDP-L-fucose in *Arabidopsis* 445
Bouchez, D. *see* Geelen, D.
Bove, J. *see* Brangeon, J.
Bovet, L. *see* Wüthrich, K.L.
Brangeon, J., Sabar, M., Gutierres, S., Combettes, B., Bove, J., Gendy, C., Chétrit, P., Colas Des Francs-Small, C., Pla, M., Vedel, F. and De Paepe, R. Defective splicing of the first *nad4* intron is associated with lack of several complex I subunits in the *Nicotiana sylvestris* NMS1 nuclear mutant 269
Brutnell, T.P. *see* Schultes, N.P.
Bush, M. *see* Karlsson, M.

Callis, J. *see* Worley, C.K.
Chétrit, P. *see* Brangeon, J.
Choe, S., Tanaka, A., Noguchi, T., Fujioka, S., Takatsuto, S., Ross, A.S., Tax, F.E., Yoshida, S. and Feldmann, K.A. Lesions in the sterol Δ^7 reductase gene of Arabidopsis cause dwarfism due to a block in brassinosteroid biosynthesis 431
Colas Des Francs-Small, C. *see* Brangeon, J.
Colcombet, J. *see* Frachisse, J.-M.
Combettes, B. *see* Brangeon, J.
Courtial, B. *see* Geelen, D.
Crosby, W.L. *see* Lui, H.
Crute, I.R. *see* Bittner-Eddy, P.D.
Czihal, A. *see* Reidt, W.

de Bruin, R. *see* Stam, M.
De Paepe, R. *see* Brangeon, J.
De Torres Zabala, M. *see* Sanchez, P.
DeLong, C. *see* Lui, H.
Dickey, L.F. *see* Petracek, M.E.
Donnison, I.S. *see* Wüthrich, K.L.
Dornelas, M.C., van Lammeren, A.A.M. and Kreis, M. *Arabidopsis thaliana* SHAGGY-related protein kinases (AtSK11 and 12) function in perianth and gynoecium development 419
Dröge, S. *see* Piedras, P.
Dry, I.B. *see* Takos, A.M.

Ellerström, M. *see* Reidt, W.
Elliott, R.C. *see* Ross, J.J.
Emura, J. *see* Ueno, Y.
Engels, C. *see* von Wirén, N.
Eymery, F. *see* Rey, P.
Ezcurra, I. *see* Reidt, W.

Feldmann, K.A. *see* Choe, S.
Fester, T. *see* Walter, M.H.
Feussner, I. *see* Maucher, H.
Fleck, O. *see* Gallego, F.
Fluhr, R. *see* Savaldi-Goldstein, S.
Fobert, P.R. *see* Lui, H.
Fowke, L.C. *see* Lui, H.
Frachisse, J.-M. *see* Geelen, D.
Frachisse, J.-M., Colcombet, J., Guern, J. and Berbier-Brygoo, H. Characterization of a nitrate-permeable channel able to mediate sustained anion efflux in hypocotyl cells from *Arabidopsis thaliana* 361
Frankard, V. *see* Vauterin, M.
Freyermuth, S.K., Bacanamwo, M. and Polacco, J.C. The soybean *Eu3* gene encodes an Ni-binding protein necessary for urease activity 53
Frommer, W.B. *see* von Wirén, N.
Fujioka, S. *see* Choe, S.

Gadella Jr, T.W.J. *see* Goedhart, J.
Gallego, F., Fleck, O., Li, A., Wyrzykowska, J. and Tinland, B. AtRAD1, a plant homologue of human and yeast nucleotide excision repair endonucleases, is involved in dark repair of UV damages and recombination 507
Geelen, D., Lurin, C., Bouchez, D., Frachisse, J.-M., Lelièvre, F., Courtial, B., Barbier-Brygoo, H. and Maurel, C. Disruption of putative anion channel gene *AtCLC-a* in *Arabidopsis* suggests a role in the regulation of nitrate content 259
Gelvin, S.B. *see* Mysore, K.S.
Gendy, C. *see* Brangeon, J.
Gillet, B. *see* Rey, P.
Gilliham, M. *see* Kiegle, E.
Gillissen, B. *see* von Wirén, N.
Goedhart, J., Hink, M.A., Visser, A.J.W.G., Bisseling, T. and Gadella Jr, T.W.J. *In vivo* fluorescence correlation microscopy (FCM) reveals accumulation and immobilization of Nod factors in root hair cell walls 109
Goldschmidt-Clermont, M. *see* Vaistij, F.E.
Grant, M. *see* Sanchez, P.
Grbić, V. and Bleecker, A.B. Axillary meristem development in *Arabidopsis thaliana* 215
Guern, J. *see* Frachisse, J.-M.
Guo, A., Salih, G. and Klessig, D.F. Activation of a diverse set of genes during the tobacco resistance response to TMV is independent of salicylic acid; induction of a subset is also ethylene independent 409
Gutierres, S. *see* Brangeon, J.
Guy, C. *see* Li, Q.-B.

Hartmann, U., Höhmann, S., Nettesheim, K., Wisman, E., Saedler, H. and Huijser, P. Molecular cloning of *SVP*: a negative regulator of the floral transition in *Arabidopsis* 351
Haskell, D. *see* Li, Q.-B.
Hasseloff, J. *see* Kiegle, E.

© 2000 Blackwell Science Ltd

Hause, B. *see* Maucher, H.
Hillmer, S. *see* Piedras, P.
Hink, M.A. *see* Goedhart, J.
Hoffmann-Thoma, G. *see* Klein, M.
Höhmann, S. *see* Hartmann, U.
Holdsworth, M.J. *see* Jones, H.D.
Holdsworth, M.J. *see* Kurup, S.
Holub, E.B. *see* Bittner-Eddy, P.D.
Hörtensteiner, S. *see* Wüthrich, K.L.
Hossain, G.S. *see* Liu, Z.
Huijser, P. *see* Hartmann, U.
Hunziker, P.E. *see* Wüthrich, K.L.

Iglesias, V.A. and Meins Jr, F. Movement of plant viruses is delayed in a β-1,3-glucanase-deficient mutant showing a reduced plasmodesmatal size exclusion limit and enhanced callose deposition 157
Imanari, E. *see* Ueno, Y.
Inami, K. *see* Ueno, Y.
Islas-Osuna, M.A. *see* Liu, Z.
Izui, K. *see* Ueno, Y.

Jacobs, M. *see* Vauterin, M.
Jeblick, W. *see* Becker, J.
Johansson, I. *see* Karlsson, M.
Jones, H.D. *see* Kurup, S.
Jones, H.D., Kurup, S., Peters, N.C.B. and Holdsworth, M.J. Identification and analysis of proteins that interact with the *Avena fatua* homologue of the maize transcription factor VIVIPAROUS 1 133
Jones, J.D.G. *see* Piedras, P.
Jost, W. *see* von Wirén, N.

Kang, H.-G. and Singh, K.B. Characterization of salicylic acid responsive, Arabidopsis Dof domain proteins: overexpression of OBP3 leads to growth defects 329
Kanyuka, K. *see* Bendahmane, A.
Karlsson, M., Johansson, I., Bush, M., McCann, M.C., Maurel, C., Larsson, C. and Kjellbom, P. An abundant TIP expressed in mature highly vacuolated cells 83
Kauss, H. *see* Becker, J.
Keller, F. *see* Sprenger, N.
Kempf, R. *see* Becker, J.
Kerckhoffs, L.H.J. *see* Ross, J.J.
Kiegle, E., Gilliham, M., Hasseloff, J. and Tester, M. Hyperpolarisation-activated calcium currents found only in cells from the elongation zone of *Arabidopsis thaliana* roots 225
Kjellbom, P. *see* Karlsson, M.
Klein, M., Martinoia, E., Hoffmann-Thoma, G. and Weissenböck, G. A membrane-potential dependent ABC-like transporter mediates the vacuolar uptake of rye flavone glucuronides: regulation of glucuronide uptake by glutathione and its conjugates 289
Klessig, D.F. *see* Guo, A.
Klos, C. *see* Seitz, B.
Kooter, J.M. *see* Stam, M.
Kranz, H., Scholz, K. and Weisshaar, B. *c-MYB* oncogene-like genes encoding three MYB repeats occur in all major plant lineages 231
Kreis, M. *see* Dornelas, M.C.
Krueger, R.W. *see* Schultes, N.P.
Kumar, C.T.R. *see* Mysore, K.S.
Kuntz, M. *see* Rey, P.
Kurup, S. *see* Jones, H.D.
Kurup, S., Jones, H.D. and Holdsworth, M.J. Interactions of the developmental regulator ABI3 with proteins identified from developing *Arabidopsis* seeds 143

Larsson, C. *see* Karlsson, M.
Lauter, F.-R. *see* von Wirén, N.
Lelièvre, F. *see* Geelen, D.
Leyser, O. *see* Worley, C.K.
Li, A. *see* Gallego, F.
Li, Q.-B., Haskell, D., Zhang, C., Sung, D.-Y. and Guy, C. Diurnal regulation of Hsp70s in leaf tissue 373
Liu, Z., Hossain, G.S., Islas-Osuna, M.A., Mitchell, D.L. and Mount, D.W. Repair of UV damage in plants by nucleotide excision repair: *Arabidopsis UVH1* DNA repair gene is a homolog of *Saccharomyces cerevisiae Rad1* 519
Lorberth, R. *see* Ritte, G.
Love, J., Scott, A.C. and Thompson, W.F. Stringent control of transgene expression in *Arabidopsis thaliana* using the Top10 promoter system 579
Lui, H., Wang, H., DeLong, C., Fowke, L.C., Crosby, W.L. and Fobert, P.R. The *Arabidopsis* Cdc2a-interacting protein ICK2 is structurally related to ICK1 and is a potent inhibitor of cyclin-dependent kinase activity *in vitro* 379
Lurin, C. *see* Geelen, D.

Martinoia, E. *see* Klein, M.
Massimino, J. *see* Rey, P.
Matsuda, N. *see* Ueda, T.
Matzk, F., Meister, A. and Schubert, I. An efficient screen for reproductive pathways using mature seeds of monocots and dicots 97
Maucher, H., Hause, B., Feussner, I., Ziegler, J. and Wasternack, C. Allene oxide synthases of barley (*Hordeum vulgare* cv. Salome): tissue specific regulation in seedling development 199
Maurel, C. *see* Geelen, D.
Maurel, C. *see* Karlsson, M.
McCann, M.C. *see* Karlsson, M.
Meins Jr, F. *see* Iglesias, V.A.
Meister, A. *see* Matzk, F.
Mitchell, D.L. *see* Liu, Z.
Mol, J.N.M. *see* Stam, M.
Monde, R.-A., Zito, F., Olive, J., Wollman, F.-A. and Stern, D.B. Post-transcriptional defects in tobacco chloroplast mutants lacking the cytochrome b_6/f complex 61
Mount, D.W. *see* Liu, Z.
Mysore, K.S., Kumar, C.T.R. and Gelvin, S.B. *Arabidopsis* ecotypes and mutants that are recalcitrant to *Agrobacterium* root transformation are susceptible to germ-line transformation 9

Nakajima, K. *see* Ueno, Y.
Nakano, A. *see* Ueda, T.
Nettesheim, K. *see* Hartmann, U.
Neubohn, B. *see* Weschke, W.
Ninnemann, O. *see* von Wirén, N.
Nishiguchi, M. *see* Sonoda, S.
Noguchi, T. *see* Choe, S.
Nuygen, T. *see* Petracek, M.E.

O'Neill, D.P. *see* Ross, J.J.
Olive, J. *see* Monde, R.-A.

Oster, U., Tanaka, R., Tanaka, A. and Rüdiger, W. Cloning and functional expression of the gene encoding the key enzyme for chlorophyll *b* biosynthesis (CAO) from *Arabidopsis thaliana* 305
Panitz, R. *see* Weschke, W.
Peltier, G. *see* Rey, P.
Peters, N.C.B. *see* Jones, H.D.
Petracek, M.E., Nuygen, T., Thompson, W.F. and Dickey, L.F. Premature termination codons destabilize ferredoxin-1 mRNA when ferredoxin-1 is translated 563
Piedras, P., Rivas, S., Dröge, S., Hillmer, S. and Jones, J.D.G. Functional, c-myc-tagged *Cf-9* resistance gene products are plasma-membrane localized and glycosylated 529
Pla, M. *see* Brangeon, J.
Polacco, J.C. *see* Freyermuth, S.K.

Querci, M. *see* Bendahmane, A.

Ramos, J. *see* Worley, C.K.
Rask, L. *see* Reidt, W.
Reidt, W., Wohlfarth, T., Ellerström, M., Czihal, A., Tewes, A., Ezcurra, I., Rask, L. and Bäumlein, H. Gene regulation during late embryogenesis: the RY motif of maturation-specific gene promoters is a direct target of the FUS3 gene product 401
Reiter, W.-D. *see* Bonin, C.P.
Rey, P., Gillet, B., Römer, S., Eymery, F., Massimino, J., Peltier, G. and Kuntz, M. Over-expression of a pepper plastid lipid-associated protein in tobacco leads to changes in plastid ultrastructure and plant development upon stress 483
Ritte, G., Lorberth, R. and Steup, M. Reversible binding of the starch-related R1 protein to the surface of transitory starch granules 387
Rivas, S. *see* Piedras, P.
Rochaix, J.-D. *see* Vaistij, F.E.
Römer, S. *see* Rey, P.
Ross, A.S. *see* Choe, S.
Ross, J.J., O'Neill, D.P., Smith, J.J., Kerckhoffs, L.H.J. and Elliott, R.C. Evidence that auxin promotes gibberellin A_1 biosynthesis in pea 547
Rouse, D. *see* Worley, C.K.
Rüdiger, W. *see* Oster, U.

Sabar, M. *see* Brangeon, J.
Saedler, H. *see* Hartmann, U.
Saedler, H. *see* Yephremov, A.
Sakakibara, H. *see* Ueno, Y.
Salih, G. *see* Guo, A.
Sanchez, P., de Torres Zabala, M. and Grant, M. AtBI-1, a plant homologue of Bax Inhibitor-1, suppresses Bax-induced cell death in yeast and is rapidly upregulated during wounding and pathogen challenge 393
Sauer, N. *see* Weschke, W.
Savaldi-Goldstein, S., Sessa, G. and Fluhr, R. The ethylene-inducible PK12 kinase mediates the phosphorylation of SR splicing factors 91
Sawers, R.J.H. *see* Schultes, N.P.
Scholz, K. *see* Kranz, H.
Schroda, M., Blöcker, D. and Beck, C.F. The *HSP70A* promoter as a tool for the improved expression of transgenes in *Chlamydomonas* 121
Schubert, I. *see* Matzk, F.
Schultes, N.P., Sawers, R.J.H., Brutnell, T.P. and Krueger, R.W. Maize *high chlorophyll fluorescent 60* mutation is caused by an *Ac* disruption of the gene encoding the chloroplast ribosomal small subunit protein 17 317
Scott, A.C. *see* Love, J.
Seitz, B., Klos, C., Wurm, M. and Tenhaken, R. Matrix polysaccharide precursors in *Arabidopsis* cell walls are synthesized by alternate pathways with organ-specific expression patterns 537
Sessa, G. *see* Savaldi-Goldstein, S.
Shiba, T. *see* Ueno, Y.
Singh, K.B. *see* Kang, H.-G.
Smith, J.J. *see* Ross, J.J.
Sonoda, S and Nishiguchi, M. Graft transmission of post-transcriptional gene silencing: target specificity for RNA degradation is transmissible between silenced and non-silenced plants, but not between silenced plants 1
Soole, K.L. *see* Takos, A.M.
Sprenger, N. and Keller, F. Allocation of raffinose family oligosaccharides to transport and storage pools in *Ajuga reptans*: the roles of two distinct galactinol synthases 249
Stam, M., de Bruin, R., van Blokland, R., van der Hoorn, R.A.L, Mol, J.N.M. and Kooter, J.M. Distinct features of post-transcriptional gene silencing by antisense transgenes in single copy and inverted T-DNA repeat loci 27
Stern, D.B. *see* Monde, R.-A.
Steup, M. *see* Ritte, G.
Strack, D. *see* Walter, M.H.
Sugiyama, T. *see* Ueno, Y.
Sung, D.-Y. *see* Li, Q.-B.

Takatsuto, S. *see* Choe, S.
Takos, A.M., Dry, I.B. and Soole, K.L. Glycosyl-phosphatidylinositol-anchor addition signals are processed in *Nicotiana tabacum* 43
Tanaka, A. *see* Choe, S.
Tanaka, A. *see* Oster, U.
Tanaka, R. *see* Oster, U.
Tax, F.E. *see* Choe, S.
Tenhaken, R. *see* Seitz, B.
Tester, M. *see* Kiegle, E.
Tewes, A. *see* Reidt, W.
Theologis, A. *see* Worley, C.K.
Thompson, W.F. *see* Love, J.
Thompson, W.F. *see* Petracek, M.E.
Tinland, B. *see* Gallego, F.

Uchimiya, H. *see* Ueda, T.
Ueda, T., Matsuda, N., Uchimiya, H. and Nakano, A. Modes of interaction between the *Arabidopsis* Rab protein, Ara4, and its putative regulator molecules revealed by a yeast expression system 341
Ueno, Y., Imanari, E., Emura, J., Yoshizawa-Kumagaye, K., Nakajima, K., Inami, K., Shiba, T., Sakakibara, H., Sugiyama, T. and Izui, K. Immunological analysis of the phosphorylation state of maize C_4-form phosphoenolpyruvate carboxylase with specific antibodies raised against a synthetic phosphorylated peptide 17

Vaistij, F.E., Goldschmidt-Clermont, M., Wostrikoff, K. and Rochaix, J.-D. Stability determinants in the chloroplast *psbB/T/H/* mRNAs of *Chlamydomonas reinhardtii* 469
van Blokland, R. *see* Stam, M.
van der Hoorn, R.A.L *see* Stam, M.
van Lammeren, A.A.M. *see* Dornelas, M.C.
Vauterin, M., Frankard, V. and Jacobs, M. Functional rescue of a bacterial *dap*A auxotroph with a plant cDNA library selects for mutant clones encoding a feedback-insensitive dihydrodipicolinate synthase 239

Vedel, F. *see* Brangeon, J.
Visser, A.J.W.G. *see* Goedhart, J.
von Wirén, N., Lauter, F.-R., Ninnemann, O., Gillissen, B., Walch-Liu, P., Engels, C., Jost, W. and Frommer, W.B. Differential regulation of three functional ammonium transporter genes by nitrogen in root hairs and by light in leaves of tomato 167
Walch-Liu, P. *see* von Wirén, N.
Walter, M.H., Fester, T. and Strack, D. Arbuscular mycorrhizal fungi induce the non-mevalonate methylerythritol phosphate pathway of isoprenoid biosynthesis correlated with accumulation of the 'yellow pigment' and other apocarotenoids 571
Wang, H. *see* Lui, H.
Wang, Q. *see* Weschke, W.
Wasternack, C. *see* Maucher, H.
Weber, H. *see* Weschke, W.
Weissenböck, G. *see* Klein, M.
Weisshaar, B. *see* Kranz, H.
Weschke, W., Panitz, R., Sauer, N., Wang, Q., Neubohn, B., Weber, H. and Wobus, U. Sucrose transport into barley seeds: molecular characterization of two transporters and implications for seed development and starch accumulation 455
Wisman, E. *see* Hartmann, U.
Wobus, U. *see* Weschke, W.
Wohlfarth, T. *see* Reidt, W.
Wollman, F.-A. *see* Monde, R.-A.
Worley, C.K., Zenser, N., Ramos, J., Rouse, D., Leyser, O., Theologis, A. and Callis, J. Degradation of Aux/IAA proteins is essential for normal auxin signalling 553
Wostrikoff, K. *see* Vaistij, F.E.
Wurm, M. *see* Seitz, B.
Wüthrich, K.L., Bovet, L., Hunziker, P.E., Donnison, I.S. and Hörtensteiner, S. Molecular cloning, functional expression and characterisation of RCC reductase involved in chlorophyll catabolism 189
Wyrzykowska, J. *see* Gallego, F.

Yanagisawa, S. Dof1 and Dof2 transcription factors are associated with expression of multiple genes involved in carbon metabolism in maize 281
Yephremov, A. and Saedler, H. Display and isolation of transposon-flanking sequences starting from genomic DNA or RNA 495
Yoshida, S. *see* Choe, S.
Yoshizawa-Kumagaye, K. *see* Ueno, Y.

Zenser, N. *see* Worley, C.K.
Zhang, C. *see* Li, Q.-B.
Ziegler, J. *see* Maucher, H.
Zito, F. *see* Monde, R.-A.

The Plant Journal (2000) **21**(6), 593–595

Subject index

aadA 121
ABC (ATP-binding cassette) transporter 289
Ac 317
Agrobacterium 9
alleles 177
allene oxide synthase 199
aluminium 225
ammonium uptake 167
anion channel 361
anion channel inhibitor 361
anion efflux 361
anion transport 259
antibodies to synthetic phosphopeptide 17
antisense 27
antisense RNA 27
apomixis 97
apoptosis 393
aquaporin 83
Ara 341
Arabidopsis 9, 53, 143, 177, 215, 259, 329, 341, 379, 393, 419, 507, 553
Arabidopsis thaliana 305, 379, 579
arginine serine rich proteins 91
arylsulfatase 121
assembly of Complex I membrane and peripheral arms 269
AtGDI 341
Aux/IAA proteins 553
auxin 329, 547, 553
Avena fatua 133
axillary meristems 215
axr3 553

barley 199, 455
Bax inhibitor 393
biosynthesis 547
branching 215
brassinolide 431

C4 gene 281
C_4 photosynthesis 17
Ca^{2+} channels 225
cab 373
carbohydrate levels 455
carbon allocation 249
carbon metabolism 281
carotenoids 571
cDNAs 199
cell death 393
cell division cycle 379
cell elongation 431
cell wall 109, 445
cell-to-cell movement 157
cellulose 157
channel 259
chitinase 311
Chlamydomonas 121
Chlamydomonas reinhardtii 469
chlorophyll breakdown 189
chlorophyllides *a* and *b* 305
chloroplast 469, 483
chloroplast import 189
circadian 373
Cladosporium fulvum 529
CLC 259
c-MYB 231
CO_2 167
compartmentation 249
complex assembly 61
cosuppression 27
cucumber 311
cyclin-dependent kinase inhibitor 379

degradation signal 553
deoxyxylulose 5-phosphate synthase 571
developmental regulation 199
diffusion 109
dihydrodipicolinate synthase 239
diurnal regulation 167
diversifying selection 177
DNA-binding protein 351
DNA methylation 1, 27
DNA repair 519
Dof protein 281
dsRNA 27

elicitation competence 311
embryo development 143
embryo dormancy 133
endoglucanase 43
endonuclease 519
ETID 495
evolution 177, 189, 231

FCS 109
Fed-1 563
feedback insensitivity 239
fibrillin 483
flavonoids 289
flow cytometric seed screen 97
flower development 419
flowering 351
fluorescence 109
functional complementation 239
FUS3 and ABI3 regulators 401

galactinol synthase 249
gene cloning 317
gene evolution 73
gene expression 281
gene families 231
gene regulation 401
gene silencing 1
germ-line transformation 9
GFP 225, 579
gibberellin 3-oxidase 547
gibberellins 547
Glomus intraradices 571
glucanase 157
glycoprotein 529
GPI-anchor 43
graft-transmission 1
GSK-3 419

© 2000 Blackwell Science Ltd

H_2O_2 311
hemicellulose 537
heterologous expression 167
high chlorophyll fluorescence 317
HSP70 121
HypB 53
hypersensitive response 73
hypocotyl 361

in situ hybridization 199
inducible promoter 579
inositol oxidation pathway 537
intermediary cells 249
inverted repeats 27

jasmonates 199

kinase 419

LAMMER kinase 91
late embryogenesis 401
LE 547
ligation-mediated PCR 495
light regulation 281, 373
light stress 483
luciferase 553
lysine biosynthesis 239

MADS-box gene 351
maize 317
major intrinsic protein 83
MEP pathway 571
mesophyll 249
MIP 83
mitochondrial encoded subunits of respiratory Complex I 269
molecular chaperone 373
monooxygenase 305
monosaccharide 445
mRNA half-life 563
mRNA stability 431, 469
mtRNA processing and translation 269
mutagenesis 239
mutants 259, 495
mycorrhiza 571

nematode 73
NER 507, 519
nickel 53
nitrate 259, 361
nitrogen nutrition 167
Nod factor 109
nonsense mediated decay 563
nuclear control of mtDNA expression 269
nuclear localization 91
nucleotide regulation 361
nucleotide sugars 445, 537

organic anions 289

parthenogenesis 97
patch clamp 225
pathogenesis-related proteins 409
patterning 419
perception 109
Peronospora 177
petA 61
petB 61
petD 61
phosphoenolpyruvate carboxylase 17
phosphorylation 91
photolyases 507
photorespiration 167
photosynthetic mutant 469
phylogeny 231, 351
Pisum sativum L. 387, 547
plant defence 529
plant nutrition 225
plants 43, 351
plasma membrane 529
plasmodesmatal trafficking 157
plastid 571
plastoglobule 483
postembryonic development 215
promoter fusions 121
proteasome activity 311
protein synthesis 373
pseudogamy 97

R2R3-MYB 231
Rab/Ypt GTPase 341
raffinose family oligosaccharides 249
recombinant enzyme 305
red chlorophyll catabolite reductase 189
regulatory phosphorylation 17
resistance 177
resistance gene 529
ribosomal protein 317
RNA processing 469
RNA stability 563
root transformation 9
RY *cis*-element 401

S. pombe 507
salicylic acid 329
SAY1 341
seed development 455
seed proteins 401
senescence 189
sexuality 97
SHAGGY 419
shoot patterning 215
signal transduction 225
signaling 409
signals 43
silencing signal 1
site-specific mutagenesis 91
size exclusion limit 157
Smith–Lemli–Opitz syndrome (SLOS) 431
Solanum tuberosum L. 387
soybean 53
spinach 373
Spinacia oleracea 83
splice site mutations 431
starch-bound proteins 387
starch granule 87
starch metabolism 387
steroid hormone 31
sugar metabolising enzymes 455

sugar transporters 455
systemic acquired resistance 311, 409
systemic spread 1

targeting 43
tetracycline 579
thylakoid 483
thymidine dimers 507
TID 495
TIP 83
tomato 529
tonoplast intrinsic protein 83
Top10 promoter 579
transactivator 579
transcription 133, 143
transcription factor 231, 281, 329, 401
transformation 61
transgene expression 121
transgene 579
translation 563
transport 289
transposition 317
transposon tagging 495

UDP-glucose dehydrogenase 537
unreduced gametes 97
unwanted selection 239
urease 53
UreG 53
UV radiation 519
UVH1 gene 519

vacuole 289
vacuum infiltration 9
vegetative growth 351
virus 73
virus resistance 1
virus speed 157

water channel 83
water deficit 483

yeast 341, 393
yeast two-hybrid 133, 143, 379

Zea mays 17
zinc finger 329

THE plant JOURNAL

VOLUME 21

Editors

Thomas Altmann
Dianna Bowles
Jeff Dangl
Jean Dénarié
Jérôme Giraudat
Harry Klee
Maarten Koornneef
Cathie Martin
Gregory Martin
Julian Schroeder
Kazuo Shinozaki
Mark Stitt

Published by Blackwell Science Ltd,
in association with the Society for Experimental Biology

© 2000 Blackwell Science Ltd and the Society for Experimental Biology. Authorization to photocopy for internal or personal use or the internal or personal use of specific clients is granted by Blackwell Science Ltd and the Society for Experimental Biology for libraries and other users registered with the Copyright Clearance Centre (CCC) Transactional Reporting Service, provided that the base fee of $15.00 per copy is paid directly to the CCC, 222 Rosewood Drive, Suite 910, Danvers, MA 01923, USA. This consent does not extend to other kinds of copying, such as copying for general distribution for advertising or promotional purposes, for creating new collective works or for resale. Special requests should be addressed to the Chief Editor.
0960-7412/00 $15.00.

Published by Blackwell Science Ltd, in association with the Society for Experimental Biology

Blackwell Science Ltd

Osney Mead, Oxford OX2 0EL, UK
25 John Street, London WC1N 2BL, UK
23 Ainslie Place, Edinburgh EH3 6AJ, UK

Blackwell Science Inc., Commerce Place, 350 Main Street, Malden, MA 02148-5018, USA
Blackwell Science Asia, 54 University Street, Carlton South, Victoria 3053, Australia
Blackwell Wissenschafts-Verlag GmbH, Kurfürstendamm 57, 10707 Berlin, Germany
Blackwell Science KK, MG-Kodenmacho Building, 7–10 Kodenmacho Nihombashi, Cho-ku, Tokyo 103-0001, Japan
Blackwell Science Ltd, 10 rue Casimir Delavigne, 75006 Paris, France

ISSN 0960-7412

Typeset and printed by Information Press Ltd, Oxford, UK.

Contents

Number 1 January 2000

v **Editorial**

1 **Graft transmission of post-transcriptional gene silencing: target specificity for RNA degradation is transmissible between silenced and non-silenced plants, but not between silenced plants.** *S. Sonoda and M. Nishiguchi*

9 ***Arabidopsis* ecotypes and mutants that are recalcitrant to *Agrobacterium* root transformation are susceptible to germ-line transformation.** *K.S. Mysore, C.T.R. Kumar and S.B. Gelvin*

17 **Immunological analysis of the phosphorylation state of maize C_4-form phosphoenolpyruvate carboxylase with specific antibodies raised against a synthetic phosphorylated peptide.** *Y. Ueno, E. Imanari, J. Emura, K. Yoshizawa-Kumagaye, K. Nakajima, K. Inami, T. Shiba, H. Sakakibara, T. Sugiyama and K. Izui*

27 **Distinct features of post-transcriptional gene silencing by antisense transgenes in single copy and inverted T-DNA repeat loci.** *M. Stam, R. de Bruin, R. van Blokland, R.A.L van der Hoorn, J.N.M. Mol and J.M. Kooter*

43 **Glycosyl-phosphatidylinositol-anchor addition signals are processed in *Nicotiana tabacum*.** *A.M. Takos, I.B. Dry and K.L. Soole*

53 **The soybean *Eu3* gene encodes an Ni-binding protein necessary for urease activity.** *S.K. Freyermuth, M. Bacanamwo and J.C. Polacco*

61 **Post-transcriptional defects in tobacco chloroplast mutants lacking the cytochrome b_6/f complex.** *R.-A. Monde, F. Zito, J. Olive, F.-A. Wollman and D.B. Stern*

73 ***Agrobacterium* transient expression system as a tool for the isolation of disease resistance genes: application to the *Rx2* locus in potato.** *A. Bendahmane, M. Querci, K. Kanyuka and D.C. Baulcombe*

83 SHORT COMMUNICATION
An abundant TIP expressed in mature highly vacuolated cells. *M. Karlsson, I. Johansson, M. Bush, M.C. McCann, C. Maurel, C. Larsson and P. Kjellbom*

91 SHORT COMMUNICATION
The ethylene-inducible PK12 kinase mediates the phosphorylation of SR splicing factors. *S. Savaldi-Goldstein, G. Sessa and R. Fluhr*

97 TECHNICAL ADVANCE
An efficient screen for reproductive pathways using mature seeds of monocots and dicots. *F. Matzk, A. Meister and I. Schubert*

109 TECHNICAL ADVANCE
***In vivo* fluorescence correlation microscopy (FCM) reveals accumulation and immobilization of Nod factors in root hair cell walls.** *J. Goedhart, M.A. Hink, A.J.W.G. Visser, T. Bisseling and T.W.J. Gadella Jr*

Number 2 January 2000

121 **The *HSP70A* promoter as a tool for the improved expression of transgenes in *Chlamydomonas*.** *M. Schroda, D. Blöcker and C.F. Beck*

133 **Identification and analysis of proteins that interact with the *Avena fatua* homologue of the maize transcription factor VIVIPAROUS 1.** *H.D. Jones, S. Kurup, N.C.B. Peters and M.J. Holdsworth*

143 **Interactions of the developmental regulator ABI3 with proteins identified from developing *Arabidopsis* seeds.** *S. Kurup, H.D. Jones and M.J. Holdsworth*

157 **Movement of plant viruses is delayed in a β-1,3-glucanase-deficient mutant showing a reduced plasmodesmatal size exclusion limit and enhanced callose deposition.** *V.A. Iglesias and F. Meins Jr*

167 **Differential regulation of three functional ammonium transporter genes by nitrogen in root hairs and by light in leaves of tomato.** *N. von Wirén, F.-R. Lauter, O. Ninnemann, B. Gillissen, P. Walch-Liu, C. Engels, W. Jost and W.B. Frommer*

177 ***RPP13* is a simple locus in *Arabidopsis thaliana* for alleles that specifiy downy mildew resistance to different avirulence determinants in *Peronospora parasitica*.** *P.D. Bittner-Eddy, I.R. Crute, E.B. Holub and J.L. Beynon*

189 **Molecular cloning, functional expression and characterisation of RCC reductase involved in chlorophyll catabolism.** *K.L. Wüthrich, L. Bovet, P.E. Hunziker, I.S. Donnison and S. Hörtensteiner*

199 **Allene oxide synthases of barley (*Hordeum vulgare* cv. Salome): tissue specific regulation in seedling development.** *H. Maucher, B. Hause, I. Feussner, J. Ziegler and C. Wasternack*

215 **Axillary meristem development in *Arabidopsis thaliana*.** *V. Grbić and A.B. Bleecker*

225 SHORT COMMUNICATION
Hyperpolarisation-activated calcium currents found only in cells from the elongation zone of *Arabidopsis thaliana* roots. *E. Kiegle, M. Gilliham, J. Hasseloff and M. Tester*

231 SHORT COMMUNICATION
***c-MYB* oncogene-like genes encoding three MYB repeats occur in all major plant lineages.** *H. Kranz, K. Scholz and B. Weisshaar*

237 **Erratum concerning Volume 20, Number 3, November 1999, pages 305–315, Fath *et al.***

Number 3 February 2000

v **Thank you**

239 **Functional rescue of a bacterial *dapA* auxotroph with a plant cDNA library selects for mutant clones encoding a feedback-insensitive dihydrodipicolinate synthase.** *M. Vauterin, V. Frankard and M. Jacobs*

249 **Allocation of raffinose oligosaccharides to transport and storage pools in *Ajuga retans*: the roles of two distinct galactinol synthases.** *N. Sprenger and F. Keller*

259 **Disruption of putative anion channel gene *AtCLC-a* in *Arabidopsis* suggests a role in the regulation of nitrate content.** *D. Geelen, C. Lurin, D. Bouchez, J.-M. Frachisse, F. Lelièvre, B. Courtial, H. Barbier-Brygoo and C. Maurel*

269 **Defective splicing of the first *nad4* intron is associated with lack of several complex I subunits in the *Nicotiana sylvestris* NMS1 nuclear mutant.** *J. Brangeon, M. Sabar, S. Gutierres, B. Combettes, J. Bove, C. Gendy, P. Chétrit, C. Colas des Francs-Small, M. Pla, F. Vedel and R. De Paepe*

281 **Dof1 and Dof2 transcription factors are associated with expression of multiple genes involved in carbon metabolism in maize.** *S. Yanagisawa*

289 **A membrane-potential dependent ABC-like transporter mediates the vacuolar uptake of rye flavone glucuronides: regulation of glucuronide uptake by glutathione and its conjugates.** *M. Klein, E. Martinoia, G. Hoffmann-Thoma and G. Weissenböck*

305 SHORT COMMUNICATION
Cloning and functional expression of the gene encoding the key enzyme for chlorophyll b biosynthesis (CAO) from *Arabidopsis thaliana*. *U. Oster, R. Tanaka, A. Tanaka and W. Rüdiger*

311 SHORT COMMUNICATION
Induction of competence for elicitation of defense responses in cucumber hypocotyls requires proteasome activity. *J. Becker, R. Kempf, W. Jeblick and H. Kauss*

Number 4 February 2000

317 **Maize *high chlorophyll fluorescent 60* mutation is caused by an *Ac* disruption of the gene encoding the chloroplast ribosomal small subunit protein 17.** *N.P. Schultes, R.J.H. Sawers, T.P. Brutnell and R.W. Krueger*

329 **Characterization of salicylic acid responsive, Arabidopsis Dof domain proteins: overexpression of OBP3 leads to growth defects.** *H.-G. Kang and K.B. Singh*

341 **Modes of interaction between the *Arabidopsis* Rab protein, Ara4, and its putative regulator molecules revealed by a yeast expression system.** *T. Ueda, N. Matsuda, H. Uchimiya and A. Nakano*

351 **Molecular cloning of *SVP*: a negative regulator of the floral transition in *Arabidopsis*.** *U. Hartmann, S. Höhmann, K. Nettesheim, E. Wisman, H. Saedler and P. Huijser*

361 **Characterization of a nitrate-permeable channel able to mediate sustained anion efflux in hypocotyl cells from *Arabidopsis thaliana*.** *J.-M. Frachisse, J. Colcombet, J. Guern and H. Berbier-Brygoo*

373 SHORT COMMUNICATION
Diurnal regulation of Hsp70s in leaf tissue. *Q.-B. Li, D. Haskell, C. Zhang, D.-Y. Sung and C. Guy*

379 SHORT COMMUNICATION
The *Arabidopsis* Cdc2a-interacting protein ICK2 is structurally related to ICK1 and is a potent inhibitor of cyclin-dependent kinase activity *in vitro*. *H. Lui, H. Wang, C. DeLong, L.C. Fowke, W.L. Crosby and P.R. Fobert*

387 SHORT COMMUNICATION
Reversible binding of the starch-related R1 protein to the surface of transitory starch granules. *G. Ritte, R. Lorberth and M. Steup*

393 SHORT COMMUNICATION
AtBI-1, a plant homologue of Bax Inhibitor-1, suppresses Bax-induced cell death in yeast and is rapidly upregulated during wounding and pathogen challenge. *P. Sanchez, M. de Torres Zabala and M. Grant*

Number 5 March 2000

401 **Gene regulation during late embryogenesis: the RY motif of maturation-specific gene promoters is a direct target of the FUS3 gene product.** *W. Reidt, T. Wohlfarth, M. Ellerström, A. Czihal, A. Tewes, I. Ezcurra, L. Rask and H. Bäumlein*

409 **Activation of a diverse set of genes during the tobacco resistance response to TMV is independent of salicylic acid; induction of a subset is also ethylene independent.** *A. Guo, G. Salih and D.F. Klessig*

419 ***Arabidopsis thaliana* SHAGGY-related protein kinases (AtSK11 and 12) function in perianth and gynoecium development.** *M.C. Dornelas, A.A.M. van Lammeren and M. Kreis*

431 **Lesions in the sterol Δ^7 reductase gene of Arabidopsis cause dwarfism due to a block in brassinosteroid biosynthesis.** *S. Choe, A. Tanaka, T. Noguchi, S. Fujioka, S. Takatsuto, A.S. Ross, F.E. Tax, S. Yoshida and K.A. Feldmann*

445 **A bifunctional epimerase-reductase acts downstream of the *MUR1* gene product and completes the *de novo* synthesis of GDP-L-fucose in *Arabidopsis*.** *C.P. Bonin and W.-D. Reiter*

455 **Sucrose transport into barley seeds: molecular characterization of two transporters and implications for seed development and starch accumulation.** *W. Weschke, R. Panitz, N. Sauer, Q. Wang, B. Neubohn, H. Weber and U. Wobus*

469 **Stability determinants in the chloroplast *psbB/T/H/* mRNAs of *Chlamydomonas reinhardtii*.** *F.E. Vaistij, M. Goldschmidt-Clermont, K. Wostrikoff and J.-D. Rochaix*

483 **Over-expression of a pepper plastid lipid-associated protein in tobacco leads to changes in plastid ultrastructure and plant development upon stress.** *P. Rey, B. Gillet, S. Römer, F. Eymery, J. Massimino, G. Peltier and M. Kuntz*

495 TECHNICAL ADVANCE
Display and isolation of transposon-flanking sequences starting from genomic DNA or RNA. *A. Yephremov and H. Saedler*

Number 6 March 2000

507 **AtRAD1, a plant homologue of human and yeast nucleotide excision repair endonucleases, is involved in dark repair of UV damages and recombination.** *F. Gallego, O. Fleck, A. Li, J. Wyrzykowska and B. Tinland*

519 **Repair of UV damage in plants by nucleotide excision repair: *Arabidopsis UVH1* DNA repair gene is a homolog of *Saccharomyces cerevisiae Rad1*.** *Z. Liu, G.S. Hossain, M.A. Islas-Osuna, D.L. Mitchell and D.W. Mount*

529 **Functional, c-myc-tagged *Cf-9* resistance gene products are plasma-membrane localized and glycosylated.** *P. Piedras, S. Rivas, S. Dröge, S. Hillmer and J.D.G. Jones*

537 **Matrix polysaccharide precursors in *Arabidopsis* cell walls are synthesized by alternate pathways with organ-specific expression patterns.** *B. Seitz, C. Klos, M. Wurm and R. Tenhaken*

547 **Evidence that auxin promotes gibberellin A_1 biosynthesis in pea.** *J.J. Ross, D.P. O'Neill, J.J. Smith, L.H.J. Kerckhoffs and R.C. Elliott*

553 **Degradation of Aux/IAA proteins is essential for normal auxin signalling.** *C.K. Worley, N. Zenser, J. Ramos, D. Rouse, O. Leyser, A. Theologis and J. Callis*

563 **Premature termination codons destabilize ferredoxin-1 mRNA when ferredoxin-1 is translated.** *M.E. Petracek, T. Nuygen, W.F. Thompson and L.F. Dickey*

571 **Arbuscular mycorrhizal fungi induce the non-mevalonate methylerythritol phospate pathway of isoprenoid biosynthesis correlated with accumulation of the 'yellow pigment' and other apocarotenoids.** *M.H. Walter, T. Fester and D. Strack*

579 TECHNICAL ADVANCE
Stringent control of transgene expression in *Arabidopsis thaliana* using the Top10 promoter system. *J. Love, A.C. Scott and W.F. Thompson*

589 **Author Index**

593 **Subject Index**